

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

Upravi 200 microgram film-coated tablets.
Upravi 400 microgram film-coated tablets.
Upravi 600 microgram film-coated tablets.
Upravi 800 microgram film-coated tablets.
Upravi 1,000 microgram film-coated tablets.
Upravi 1,200 microgram film-coated tablets.
Upravi 1,400 microgram film-coated tablets.
Upravi 1,600 microgram film-coated tablets.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Upravi 200 microgram film-coated tablets

Each film-coated tablet contains 200 micrograms of selexipag.

Upravi 400 microgram film-coated tablets

Each film-coated tablet contains 400 micrograms of selexipag.

Upravi 600 microgram film-coated tablets

Each film-coated tablet contains 600 micrograms of selexipag.

Upravi 800 microgram film-coated tablets

Each film-coated tablet contains 800 micrograms of selexipag.

Upravi 1,000 microgram film-coated tablets

Each film-coated tablet contains 1,000 micrograms of selexipag.

Upravi 1,200 microgram film-coated tablets

Each film-coated tablet contains 1,200 micrograms of selexipag.

Upravi 1,400 microgram film-coated tablets

Each film-coated tablet contains 1,400 micrograms of selexipag.

Upravi 1,600 microgram film-coated tablets

Each film-coated tablet contains 1,600 micrograms of selexipag.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet

Upravi 200 microgram film-coated tablets

Round, light-yellow, film-coated tablets with “2” debossed on one side.

Upravi 400 microgram film-coated tablets

Round, red, film-coated tablets with “4” debossed on one side.

Upravi 600 microgram film-coated tablets

Round, light-violet, film-coated tablets with “6” debossed on one side.

Uptravi 800 microgram film-coated tablets

Round, green, film-coated tablets with “8” debossed on one side.

Uptravi 1,000 microgram film-coated tablets

Round, orange, film-coated tablets with “10” debossed on one side.

Uptravi 1,200 microgram film-coated tablets

Round, dark-violet, film-coated tablets with “12” debossed on one side.

Uptravi 1,400 microgram film-coated tablets

Round, dark-yellow, film-coated tablets with “14” debossed on one side.

Uptravi 1,600 microgram film-coated tablets

Round, brown, film-coated tablets with “16” debossed on one side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Uptravi is indicated for the long-term treatment of pulmonary arterial hypertension (PAH) in adult patients with WHO functional class (FC) II–III, either as combination therapy in patients insufficiently controlled with an endothelin receptor antagonist (ERA) and/or a phosphodiesterase type 5 (PDE-5) inhibitor, or as monotherapy in patients who are not candidates for these therapies.

Efficacy has been shown in a PAH population including idiopathic and heritable PAH, PAH associated with connective tissue disorders, and PAH associated with corrected simple congenital heart disease (see section 5.1).

4.2 Posology and method of administration

Treatment should only be initiated and monitored by a physician experienced in the treatment of PAH.

Posology

Individualised dose titration

Each patient should be up-titrated to the highest individually tolerated dose, which can range from 200 micrograms given twice daily to 1,600 micrograms given twice daily (individualised maintenance dose).

The recommended starting dose is 200 micrograms given twice daily, approximately 12 hours apart. The dose is increased in increments of 200 micrograms given twice daily, usually at weekly intervals. At the beginning of treatment and at each up-titration step it is recommended to take the first dose in the evening. During dose titration some adverse reactions, reflecting the mode of action of Uptravi (such as headache, diarrhoea, nausea and vomiting, jaw pain, myalgia, pain in extremity, arthralgia, and flushing), may occur. They are usually transient or manageable with symptomatic treatment (see section 4.8). However, if a patient reaches a dose that cannot be tolerated, the dose should be reduced to the previous dose level.

In patients in whom up-titration was limited by reasons other than adverse reactions reflecting the mode of action of Uptravi, a second attempt to continue up-titration to the highest individually tolerated dose up to a maximum dose of 1,600 micrograms twice daily may be considered.

Individualised maintenance dose

The highest tolerated dose reached during dose titration should be maintained. If the therapy over time is less tolerated at a given dose, symptomatic treatment and/or a dose reduction to the next lower dose

should be considered.

Interruptions and discontinuations

If a dose is missed, it should be taken as soon as possible. The missed dose should not be taken if the next scheduled dose is within approximately 6 hours.

If treatment is missed for 3 days or more, Uptravi should be restarted at a lower dose and then up-titrated.

There is limited experience with abrupt discontinuation of Uptravi in patients with PAH. No evidence for acute rebound has been observed.

However, if the decision to withdraw Uptravi is taken, it should be done gradually while an alternative therapy is introduced.

Elderly (≥ 65 years)

No adjustment to the dose regimen is needed in elderly people (see section 5.2). There is limited clinical experience in patients over the age of 75 years, therefore Uptravi should be used with caution in this population (see section 4.4).

Hepatic impairment

Uptravi should not be administered in patients with severe liver impairment (Child-Pugh class C; see section 4.4). For patients with moderate hepatic impairment (Child-Pugh class B), the starting dose of Uptravi should be 200 micrograms once daily, and increased at weekly intervals by increments of 200 micrograms given once daily until adverse reactions, reflecting the mode of action of selexipag, that cannot be tolerated or medically managed are experienced. No adjustment to the dose regimen is needed in patients with mild hepatic impairment (Child-Pugh class A).

Renal impairment

No adjustment to the dose regimen is needed in patients with mild or moderate renal impairment. No change in starting dose is required in patients with severe renal impairment (estimated glomerular filtration rate [eGFR] < 30 mL/min/1.73 m²); dose titration should be done with caution in these patients (see section 4.4).

Paediatric population (< 18 years)

The safety and efficacy of Uptravi in children aged 0 to less than 18 years have not yet been established. No data are available. Administration of selexipag in the paediatric population is not recommended. Animal studies indicated an increased risk of intussusception, but the clinical relevance of these findings is unknown (see section 5.3).

Method of administration

Oral use.

The film-coated tablets are to be taken orally in the morning and in the evening. To improve tolerability, it is recommended to take Uptravi with food and, at the beginning of each up-titration phase, to take the first increased dose in the evening.

The tablets should not be split, crushed or chewed, and are to be swallowed with water.

Patients who have poor vision or are blind must be instructed to get assistance from another person when taking Uptravi during the titration period.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Severe coronary heart disease or unstable angina.

- Myocardial infarction within the last 6 months.
- Decompensated cardiac failure if not under close medical supervision.
- Severe arrhythmias.
- Cerebrovascular events (e.g., transient ischaemic attack, stroke) within the last 3 months.
- Congenital or acquired valvular defects with clinically relevant myocardial function disorders not related to pulmonary hypertension.
- Concomitant use of strong inhibitors of CYP2C8 (e.g., gemfibrozil; see section 4.5).

4.4 Special warnings and precautions for use

Hypotension

Upravi has vasodilatory properties that may result in lowering of blood pressure. Before prescribing Upravi, physicians should carefully consider whether patients with certain underlying conditions could be adversely affected by vasodilatory effects (e.g., patients on antihypertensive therapy or with resting hypotension, hypovolaemia, severe left ventricular outflow obstruction or autonomic dysfunction).

Hyperthyroidism

Hyperthyroidism has been observed with Upravi. Thyroid function tests are recommended as clinically indicated in the presence of signs or symptoms of hyperthyroidism.

Pulmonary veno-occlusive disease

Cases of pulmonary oedema have been reported with vasodilators (mainly prostacyclins) when used in patients with pulmonary veno-occlusive disease. Consequently, if signs of pulmonary oedema occur when Upravi is administered in patients with PAH, the possibility of pulmonary veno-occlusive disease should be considered. If confirmed, treatment with Upravi should be discontinued.

Moderate inhibitors of CYP2C8

Concomitant administration of selexipag with moderate inhibitors of CYP2C8 (e.g. clopidogrel, deferasirox, teriflunomide) may increase the exposure to selexipag and its main active metabolite. An adjustment of the dose of selexipag should be considered in case a moderate inhibitor of CYP2C8 is coadministered or discontinued.

Elderly (≥ 65 years)

There is limited clinical experience with selexipag in patients over the age of 75 years, therefore Upravi should be used with caution in this population (see section 4.2).

Hepatic impairment

There is no clinical experience with selexipag in patients with severe liver impairment (Child-Pugh class C), therefore Upravi should not be administered in these patients. The exposure to selexipag and its active metabolite is increased in subjects with moderate hepatic impairment (Child-Pugh class B; see section 5.2). In patients with moderate hepatic impairment, Upravi should be dosed once daily (see section 4.2).

Renal impairment

In patients with severe renal impairment (eGFR < 30 mL/min/1.73 m²), caution should be exercised during dose titration. There is no experience with Upravi in patients undergoing dialysis (see section 5.2), therefore Upravi should not be used in these patients.

Women of childbearing potential

Women of childbearing potential should practise effective contraception while taking selexipag.

4.5 Interaction with other medicinal products and other forms of interaction

Effect of other medicinal products on selexipag

Selexipag is hydrolysed to its active metabolite by carboxylesterases (see section 5.2). Selexipag and its active metabolite both undergo oxidative metabolism mainly by CYP2C8 and to a smaller extent by CYP3A4. The glucuronidation of the active metabolite is catalysed by UGT1A3 and UGT2B7. Selexipag and its active metabolite are substrates of OATP1B1 and OATP1B3. Selexipag is a weak substrate of the P-gp efflux pump. The active metabolite is a weak substrate of the breast cancer resistance protein (BCRP).

The pharmacokinetics of selexipag and its active metabolite are not affected by warfarin.

Inhibitors of CYP2C8

In the presence of 600 mg gemfibrozil, twice a day, a strong inhibitor of CYP2C8, exposure to selexipag increased approximately 2-fold, whereas exposure to the active metabolite, the major contributor to efficacy, increased approximately 11-fold. Concomitant administration of Upravi with strong inhibitors of CYP2C8 (e.g., gemfibrozil) is contraindicated (see section 4.3).

The effect of moderate inhibitors of CYP2C8 (e.g., clopidogrel, deferasirox, teriflunomide) on the exposure to selexipag and its active metabolite has not been studied. An adjustment of the dose of Upravi should be considered in case a moderate inhibitor of CYP2C8 is coadministered or discontinued. A potential pharmacokinetic interaction with moderate inhibitors of CYP2C8 cannot be excluded (see section 4.4).

Inducers of CYP2C8

In the presence of 600 mg rifampicin, once a day, an inducer of CYP2C8 (and UGT enzymes), the exposure to selexipag did not change, whereas exposure to the active metabolite was reduced by half. Dose adjustment of selexipag may be required with concomitant administration of inducers of CYP2C8 (e.g., rifampicin, carbamazepine, phenytoin).

Inhibitors of UGT1A3 and UGT2B7

The effect of strong inhibitors of UGT1A3 and UGT2B7 (valproic acid, probenecid, and fluconazole) on the exposure to selexipag and its active metabolite has not been studied. Caution is required when administering these medicinal products concomitantly with Upravi. A potential pharmacokinetic interaction with strong inhibitors of UGT1A3 and UGT2B7 cannot be excluded.

Inhibitors and inducers of CYP3A4

In the presence of 400/100 mg lopinavir/ritonavir twice daily, a strong CYP3A4 inhibitor, exposure to selexipag increased approximately 2-fold, whereas the exposure to the active metabolite of selexipag did not change. Given the 37-fold higher potency of the active metabolite, this effect is not clinically relevant. Since a strong inhibitor of CYP3A4 did not affect the pharmacokinetics of the active metabolite, indicating that the CYP3A4 pathway is not important in the elimination of the active metabolite, no effect of inducers of CYP3A4 on the pharmacokinetics of the active metabolite is expected.

PAH-specific therapies

In the Phase 3 placebo-controlled trial in patients with PAH, the use of selexipag in combination with both an ERA and a PDE-5 inhibitor resulted in a 30% lower exposure to the active metabolite.

Transporter inhibitors (lopinavir/ritonavir)

In the presence of 400/100 mg lopinavir/ritonavir twice daily, a strong OATP (OATP1B1 and OATP1B3) and P-gp inhibitor, exposure to selexipag increased approximately 2-fold, whereas the

exposure to the active metabolite of selexipag did not change. Given that the majority of the pharmacological effect is driven by the active metabolite, this effect is not clinically relevant.

Effect of selexipag on other medicinal products

Selexipag and its active metabolite do not inhibit cytochrome P450 enzymes at clinically relevant concentrations. Selexipag and its active metabolite do not inhibit transport proteins. Selexipag and its active metabolite are not expected to induce cytochrome P450 enzymes in the liver and kidney at clinically relevant concentrations. *In vitro* data indicate that selexipag could be an inducer of both CYP3A4 and CYP2C9 in the intestine.

Anticoagulants or inhibitors of platelet aggregation

Selexipag is an inhibitor of platelet aggregation *in vitro*. In the Phase 3 placebo-controlled study in patients with PAH, no increased risk of bleeding was detected with selexipag compared to placebo, including when selexipag was administered with anticoagulants (such as heparin, coumarin-type anticoagulants) or inhibitors of platelet aggregation. In a study in healthy subjects, selexipag (400 micrograms twice daily) did not alter the exposure to S-warfarin (CYP2C9 substrate) or R-warfarin (CYP3A4 substrate) after a single dose of 20 mg warfarin. Selexipag did not influence the pharmacodynamic effect of warfarin on the international normalised ratio.

Hormonal contraceptives

Specific drug-drug interaction studies with hormonal contraceptives have not been conducted. Since selexipag did not affect the exposure to the CYP3A4 substrate R-warfarin or to the CYP2C9 substrate S-warfarin, reduced efficacy of hormonal contraceptives is not expected.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Women of childbearing potential should practise effective contraception while taking selexipag.

Pregnancy

There are no data from the use of selexipag in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. Selexipag and its main metabolite showed 20- to 80-times lower prostacyclin (IP) receptor potency *in vitro* for animal species used in reproductive toxicity testing compared to humans. Therefore, safety margins for potential IP receptor-mediated effects on reproduction are accordingly lower than for non-IP-related effects (see section 5.3).

Uptravi is not recommended during pregnancy and in women of childbearing potential not using contraception.

Breast-feeding

It is unknown whether selexipag or its metabolites are excreted in human milk. In rats, selexipag or its metabolites are excreted in milk (see section 5.3). A risk to the suckling child cannot be excluded. Uptravi should not be used during breast-feeding.

Fertility

There are no clinical data available. In rat studies, selexipag at high doses caused transient disturbances in oestrus cycles that did not affect fertility (see section 5.3). The relevance for humans is not known.

4.7 Effects on ability to drive and use machines

Uptravi has minor influence on the ability to drive and use machines. The clinical status of the patient and the adverse reaction profile of selexipag (such as headache or hypotension) should be kept in mind when considering the patient's ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

The most commonly reported adverse reactions are headache, diarrhoea, nausea and vomiting, jaw pain, myalgia, pain in extremity, arthralgia, and flushing. These reactions are more frequent during the up-titration phase. The majority of these reactions are of mild to moderate intensity.

Tabulated list of adverse reactions

The safety of selexipag has been evaluated in a long-term, Phase 3 placebo-controlled study enrolling 1,156 patients with symptomatic PAH. The mean treatment duration was 76.4 weeks (median 70.7 weeks) for patients receiving selexipag versus 71.2 weeks (median 63.7 weeks) for patients on placebo. The exposure to selexipag was up to 4.2 years.

Adverse reactions obtained from the pivotal clinical study are tabulated below. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

System organ class	<u>Very common</u> (≥ 1/10)	<u>Common</u> (≥ 1/100 to < 1/10)	<u>Uncommon</u> (≥ 1/1,000 to < 1/100)
Blood and lymphatic disorders		Anaemia Haemoglobin decreased	
Endocrine disorders		Hyperthyroidism Thyroid-stimulating Hormone decreased (see section 4.4)	
Metabolism and nutrition disorders		Decreased appetite Weight decrease	
Nervous system disorders	Headache*		
Cardiac disorders			Sinus tachycardia (see section 4.4)
Vascular disorders	Flushing*	Hypotension (see section 4.4)	
Respiratory, thoracic and mediastinal disorders	Nasopharyngitis (of non-infectious origin)	Nasal congestion	
Gastro-intestinal disorders	Diarrhoea* Vomiting * Nausea*	Abdominal pain	
Skin and subcutaneous tissue disorders		Rash Urticaria Erythema	
Musculoskeletal and connective tissue disorders	Jaw pain* Myalgia* Arthralgia* Pain in extremity*		
General disorders and administration site conditions		Pain	

* See section Description of selected adverse reactions.

Description of selected adverse reactions

Pharmacological effects associated with titration and maintenance treatment

Adverse reactions associated with the mode of action of selexipag have been observed frequently, in particular during the phase of individualised dose titration, and are tabulated below:

Prostacyclin-like associated adverse reactions	Titration		Maintenance	
	Selexipag	Placebo	Selexipag	Placebo
Headache	64%	28%	40%	20%
Diarrhoea	36%	12%	30%	13%
Nausea	29%	13%	20%	10%
Pain in jaw	26%	4%	21%	4%
Myalgia	15%	5%	9%	3%
Pain in extremity	14%	5%	13%	6%
Vomiting	14%	4%	8%	6%
Flushing	11%	4%	10%	3%
Arthralgia	7%	5%	9%	5%

These effects are usually transient or manageable with symptomatic treatment. 7.5% of patients on selexipag discontinued treatment due to these adverse reactions. The approximate rate of adverse reactions that were serious was 2.3% in the selexipag group and 0.5% in the placebo group. In clinical practice, gastro-intestinal events have been observed to respond to anti-diarrhoeal, anti-emetic, and anti-nauseant medicinal products and/or medicinal products for functional gastro-intestinal disorders. Pain-associated events have frequently been treated with analgesics (such as paracetamol).

Haemoglobin decrease

In a Phase 3 placebo-controlled study in patients with PAH, mean absolute changes in haemoglobin at regular visits compared to baseline ranged from -0.34 to -0.02 g/dL in the selexipag group compared to -0.05 to 0.25 g/dL in the placebo group. A decrease from baseline in haemoglobin concentration to below 10 g/dL was reported in 8.6% of selexipag-treated patients and 5.0% of placebo-treated patients.

Thyroid function tests

In a Phase 3 placebo-controlled study in patients with PAH, hyperthyroidism was reported for 1.6% of patients in the selexipag group, compared to no case in the placebo group (see section 4.4). A reduction (up to -0.3 MU/L from a baseline median of 2.5 MU/L) in median thyroid-stimulating hormone was observed at most visits in the selexipag group. In the placebo group, little change in median values was apparent. There were no mean changes in triiodothyronine or thyroxine in either group.

Increase in heart rate

In the Phase 3 placebo-controlled study in patients with PAH, a transient increase in mean heart rate of 3–4 bpm at 2–4 hours post-dose was observed. Electrocardiogram investigations showed sinus tachycardia in 11.3% of patients in the selexipag group compared to 8.8% in the placebo group (see also sections 4.4 and 5.1).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed below.

United Kingdom

Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard

Ireland

HPRA Pharmacovigilance
Earlsfort Terrace
IRL - Dublin 2
Tel: +353 1 6764971
Fax: +353 1 6762517
Website: www.hpra.ie
e-mail: medsafety@hpra.ie

Malta

ADR Reporting
Website: www.medicinesauthority.gov.mt/adrportal

4.9 Overdose

Isolated cases of overdose up to 3,200 micrograms were reported. Mild, transient nausea was the only reported consequence. In the event of overdose, supportive measures must be taken as required. Dialysis is unlikely to be effective because selexipag and its active metabolite are highly protein bound.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antithrombotic agents, platelet aggregation inhibitors excluding heparin, ATC code: B01AC27

Mechanism of action

Selexipag is a selective IP receptor agonist distinct from prostacyclin and its analogues. Selexipag is hydrolysed by carboxylesterases to yield its active metabolite, which is approximately 37-fold more potent than selexipag. Selexipag and the active metabolite are high-affinity IP receptor agonists with a high selectivity for the IP receptor versus other prostanoid receptors (EP₁–EP₄, DP, FP, and TP). Selectivity against EP₁, EP₃, FP, and TP is important because these are well-described contractile receptors in the gastro-intestinal tract and blood vessels. Selectivity against EP₂, EP₄, and DP₁ is important because these receptors mediate immune depressive effects.

Stimulation of the IP receptor by selexipag and the active metabolite leads to vasodilatory as well as anti-proliferative and anti-fibrotic effects. Selexipag prevents cardiac and pulmonary remodelling in a rat model of PAH and causes proportional decreases in pulmonary and peripheral pressures, indicating that peripheral vasodilation reflects pulmonary pharmacodynamic efficacy. Selexipag does not cause IP receptor desensitisation *in vitro* nor tachyphylaxis in a rat model.

Pharmacodynamic effects

Cardiac electrophysiology

In a thorough QT study in healthy subjects, repeated doses of 800 and 1,600 micrograms of selexipag twice daily did not show an effect on cardiac repolarisation (QT_c interval) or conduction (PR and QRS intervals) and had a mild accelerating effect on heart rate (the placebo-corrected, baseline-adjusted increase in heart rate reached 6–7 bpm at 1.5 to 3 h after dosing with 800 micrograms selexipag and 9–10 bpm at the same timepoints after 1,600 micrograms selexipag).

Coagulation factors

In Phase 1 and 2 studies a slight decrease in plasma levels of von Willebrand factor (vWF) was observed with selexipag; the vWF values remained above the lower limit of the normal range.

Pulmonary haemodynamics

A Phase 2 double-blind, placebo-controlled clinical study assessed haemodynamic variables after 17 weeks of treatment in patients with PAH WHO FC II–III and concomitantly receiving ERAs and/or PDE-5 inhibitors. Patients titrating selexipag to an individually tolerated dose (200 micrograms twice daily increments up to 800 micrograms twice daily; N = 33) achieved a statistically significant mean reduction in pulmonary vascular resistance of 30.3% (95% confidence interval [CI]: -44.7%, -12.2%; $p = 0.0045$) and an increase in cardiac index (mean treatment effect) of 0.48 L/min/m² (95% CI: 0.13, 0.83) compared to placebo (N = 10).

Clinical efficacy and safety

Efficacy in patients with PAH

The effect of selexipag on progression of PAH was demonstrated in a multi-centre, long-term (maximum duration of exposure approximately 4.2 years), double-blind, placebo-controlled, parallel-group, event-driven Phase 3 study in 1,156 patients with symptomatic (WHO FC I–IV) PAH. Patients were randomised to either placebo (N = 582) or selexipag (N = 574) twice daily. The dose was increased at weekly intervals by increments of 200 micrograms given twice daily to determine the individualised maintenance dose (200–1,600 micrograms twice daily).

The primary study endpoint was the time to first occurrence of a morbidity or mortality event up to end of treatment, defined as a composite of death (all causes); or hospitalisation for PAH; or progression of PAH resulting in need for lung transplantation or balloon atrial septostomy; or initiation of parenteral prostanoid therapy or chronic oxygen therapy; or other disease-progression events (patients in WHO FC II or III at baseline) confirmed by a decrease in 6-minute walk distance (6MWD) from baseline ($\geq 15\%$) and worsening of WHO FC or (patients in WHO FC III or IV at baseline) confirmed by a decrease in 6MWD from baseline ($\geq 15\%$) and need for additional PAH-specific therapy.

All events were confirmed by an independent adjudication committee, blinded to treatment allocation.

The mean age was 48.1 years (range 18–80 years of age), with the majority of subjects being Caucasian (65.0%) and female (79.8%). 17.9% of patients were ≥ 65 and 1.1% ≥ 75 years of age. Approximately 1%, 46%, 53%, and 1% of patients were in WHO FC I, II, III and IV, respectively, at baseline.


Idiopathic or heritable PAH was the most common aetiology in the study population (58%) followed by PAH due to connective tissue disorders (29%), PAH associated with simple corrected congenital heart disease (10%), and PAH associated with other aetiologies (drugs and toxins [2%] and HIV [1%]).

At baseline, the majority of enrolled patients (80%) were being treated with a stable dose of a specific therapy for PAH, either an ERA (15%) or a PDE-5 inhibitor (32%) or both an ERA and a PDE-5 inhibitor (33%).

The overall median double-blind treatment duration was 63.7 weeks for the placebo group and 70.7 weeks for the selexipag group. 23% of patients on selexipag achieved maintenance doses in the range of 200–400 micrograms, 31% achieved doses in the range of 600–1,000 micrograms, and 43% achieved doses in the range of 1,200–1,600 micrograms.

Treatment with selexipag 200–1,600 micrograms twice daily resulted in a 40% reduction (hazard ratio [HR] 0.60; 99% CI: 0.46, 0.78; one-sided log-rank p value < 0.0001) of the occurrence of morbidity or mortality events up to 7 days after last dose compared to placebo (Figure 1). The beneficial effect of selexipag was primarily attributable to a reduction in hospitalisation for PAH and a reduction in other disease-progression events (Table 1).

Figure 1 Kaplan-Meier estimates of the first morbidity-mortality event



UPTRAVI patients:	at risk	574	455	361	246	171	101	40
Placebo patients:	at risk	582	433	347	220	149	88	28

Table 1 Summary of outcome events

Endpoints & statistics	Patients with an event		Treatment comparison: selexipag vs placebo			
	Placebo (N=582)	Selexipag (N=574)	Absolute risk reduction	Relative risk reduction (99% CI)	HR (99% CI)	p-value
Morbidity-mortality event^a	58.2%	41.7%	16.5%	40% (22%; 54%)	0.60 (0.46; 0.78)	< 0.0001
Hospitalisation due to PAH^b n (%)	109 (18.7%)	78 (13.6%)	5.1%	33% (2%; 54%)	0.67 (0.46; 0.98)	0.04
Disease progression^b n (%)	100 (17.2%)	38 (6.6%)	10.6%	64% (41%; 78%)	0.36 (0.22; 0.59)	< 0.0001

i.v./s.c. Prostanoid initiation or oxygen therapy^{b,c} n (%)	15 (2.6%)	11 (1.9%)	0.7%	32% (-90%; 76%)	0.68 (0.24; 1.90)	0.53
Death up to EOT + 7 days^d n (%)	37 (6.4%)	46 (8.0%)	-1.7%	-17% (-107%; 34%)	1.17 (0.66, 2.07)	0.77
Death up to study closure^d n (%)	105 (18.0%)	100 (17.4%)	0.6%	3% (-39%; 32%)	0.97 (0.68; 1.39)	0.42

CI = confidence interval; EOT = end of treatment; HR = hazard ratio; i.v. = intravenous; PAH = pulmonary arterial hypertension; s.c. = subcutaneous.

(a) % of patients with an event at 36 months = $100 \times (1 - \text{Kaplan-Meier estimate})$; hazard ratio estimated using Cox's proportional hazard model; unstratified one-sided log-rank p-value

(b) % of patients with an event as part of the primary endpoint up to EOT + 7 days; hazard ratio estimated using Aalen Johansen method; 2-sided p-value using Gray's test

(c) Includes 'Need for lung transplantation or atrial septostomy' (1 patient on selexipag and 2 on placebo)

(d) % of patients with an event up to EOT + 7 days or up to study closure; hazard ratio estimated using Cox's proportional hazard model; unstratified one-sided log-rank p-value

The numerical increase in deaths up to end of treatment + 7 days but not up to study closure was further investigated by mathematical modelling, showing that the imbalance in deaths is consistent with the assumption of a neutral effect on PAH mortality and reduction of non-fatal events.

The observed effect of selexipag versus placebo on the primary endpoint was consistent across individualised maintenance dose as shown by the hazard ratio for the three pre-defined categories (0.60 for 200–400 micrograms twice daily, 0.53 for 600–1,000 micrograms twice daily, and 0.64 for 1,200–1,600 micrograms twice daily), which was consistent with the overall treatment effect (0.60).

The efficacy of selexipag on the primary endpoint was consistent across subgroups of age, sex, race, aetiology, geographical region, WHO FC, and as monotherapy or in combination with an ERA or a PDE-5 inhibitor or triple combination with both an ERA and a PDE-5 inhibitor.

Time to PAH-related death or hospitalisation for PAH was assessed as a secondary endpoint. The risk of an event for this endpoint was reduced by 30% in patients receiving selexipag compared to placebo (HR 0.70, 99% CI: 0.50, 0.98; one-sided log-rank $p = 0.0031$). The percentages of patients with an event at Month 36 were 28.9% and 41.3% in the selexipag and placebo groups, respectively, with an absolute risk reduction of 12.4%.

The number of patients who experienced, as a first event, death due to PAH or hospitalisation for PAH up to end of treatment was 102 (17.8%) in the selexipag group and 137 (23.5%) in the placebo group. Death due to PAH as a component of the endpoint was observed in 16 (2.8%) patients on selexipag and 14 (2.4%) on placebo. Hospitalisation for PAH was observed in 86 (15.0%) patients on selexipag and 123 (21.1%) patients on placebo. Selexipag reduced the risk of hospitalisation for PAH as a first outcome event compared to placebo (HR 0.67, 99% CI: 0.46, 0.98; one-sided log-rank $p = 0.04$).

The total number of deaths of all causes up to study closure was 100 (17.4%) for the selexipag group and 105 (18.0%) for the placebo group (HR 0.97, 99% CI: 0.68, 1.39). The number of deaths due to PAH up to study closure was 70 (12.2%) for the selexipag group and 83 (14.3%) for the placebo group.

Symptomatic endpoints

Exercise capacity was evaluated as a secondary endpoint. Median 6MWD at baseline was 376 m (range: 90–482 m) and 369 m (range: 50–515 m) in selexipag patients and placebo patients, respectively. Treatment with selexipag resulted in a placebo-corrected median effect on 6MWD measured at trough (i.e., approximately 12 h post-dose) of 12 m at Week 26 (99% CI: 1, 24 m;

one-sided p value = 0.0027). In patients without concurrent PAH-specific therapy, the placebo-corrected treatment effect measured at trough was 34 m (99% CI: 10, 63 m).

Quality of life was assessed in a subset of patients in the GRIPHON study using the Cambridge Pulmonary Hypertension Outcome Review (CAMPHOR) questionnaire. There was no significant treatment effect from baseline to Week 26.

Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Uptravi in one or more subsets of the paediatric population for treatment of pulmonary hypertension, (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

The pharmacokinetics of selexipag and its active metabolite have been studied primarily in healthy subjects. The pharmacokinetics of selexipag and the active metabolite, both after single- and multiple-dose administration, were dose-proportional up to a single dose of 800 micrograms and multiple doses of up to 1,800 micrograms twice daily. After multiple-dose administration, steady state conditions of selexipag and the active metabolite were reached within 3 days. No accumulation in plasma, either of parent compound or active metabolite, occurred after multiple-dose administration.

In healthy subjects, inter-subject variability in exposure (area under the curve over a dosing interval) at steady state was 43% and 39% for selexipag and the active metabolite, respectively. Intra-subject variability in exposure was 24% and 19% for selexipag and the active metabolite, respectively.

Exposure to selexipag and the active metabolite at steady state in PAH patients and healthy subjects was similar. The pharmacokinetics of selexipag and the active metabolite in PAH patients were not influenced by the severity of the disease and did not change with time.

Absorption

Selexipag is rapidly absorbed and is hydrolysed by carboxylesterases to its active metabolite.

Maximum observed plasma concentrations of selexipag and its active metabolite after oral administration are reached within 1–3 h and 3–4 h, respectively.

The absolute bioavailability of selexipag in humans is approximately 49%. This is most probably due to a first-pass effect of selexipag, as plasma concentrations of the active metabolite are similar after the same oral and intravenous dose administration.

In the presence of food, the exposure to selexipag after a single dose of 400 micrograms was increased by 10% in Caucasian subjects and decreased by 15% in Japanese subjects, whereas exposure to the active metabolite was decreased by 27% (Caucasian subjects) and 12% (Japanese subjects). More subjects reported adverse events after administration in the fasted than in the fed state.

Distribution

Selexipag and its active metabolite are highly bound to plasma proteins (approximately 99% in total and to the same extent to albumin and alpha1-acid glycoprotein). The volume of distribution of selexipag at steady state is 11.7 L.

Biotransformation

Selexipag is hydrolysed to its active metabolite in the liver and in the intestine by carboxylesterases. Oxidative metabolism catalysed mainly by CYP2C8 and to a smaller extent by CYP3A4 leads to the formation of hydroxylated and dealkylated products. UGT1A3 and UGT2B7 are involved in the

glucuronidation of the active metabolite. Except for the active metabolite, none of the circulating metabolites in human plasma exceed 3% of the total drug-related material. Both in healthy subjects and PAH patients, after oral administration, exposure at steady state to the active metabolite is approximately 3- to 4-fold higher than to the parent compound.

Elimination

Elimination of selexipag is predominantly via metabolism with a mean terminal half-life of 0.8–2.5 h. The active metabolite has a half-life of 6.2–13.5 h. The total body clearance of selexipag is 17.9 L/h. Excretion in healthy subjects was complete 5 days after administration and occurred primarily via faeces (accounting for 93% of the administered dose) compared to 12% in urine.

Special populations

No clinically relevant effects of sex, race, age, or body weight on the pharmacokinetics of selexipag and its active metabolite have been observed in healthy subjects or PAH patients.

Renal impairment

A 1.4- to 1.7-fold increase in exposure (maximum plasma concentration and area under the plasma concentration-time curve) to selexipag and its active metabolite was observed in subjects with severe renal impairment (eGFR < 30 mL/min/1.73 m²).

Hepatic impairment

In subjects with mild (Child-Pugh class A) or moderate (Child-Pugh class B) hepatic impairment, exposure to selexipag was 2- and 4-fold higher, respectively, when compared to healthy subjects. Exposure to the active metabolite remained almost unchanged in subjects with mild hepatic impairment and was doubled in subjects with moderate hepatic impairment. Only two subjects with severe (Child-Pugh class C) hepatic impairment were dosed with selexipag. Exposure to selexipag and its active metabolite in these two subjects was similar to that in subjects with moderate (Child-Pugh class B) hepatic impairment.

Based on modelling and simulation data from a study in subjects with hepatic impairment, the exposure to selexipag at steady state in subjects with moderate hepatic impairment (Child-Pugh class B) after a once-daily regimen is predicted to be approximately 2-fold higher than that in healthy subjects during a twice-daily regimen. The exposure to the active metabolite at steady state in these patients during a once-daily regimen is predicted to be similar to that in healthy subjects during a twice-daily regimen. Subjects with severe hepatic impairment (Child-Pugh class C) showed similar predicted exposure at steady state as subjects with moderate hepatic impairment during a once-daily regimen.

5.3 Preclinical safety data

In the repeated-dose toxicity studies in rodents, strong blood pressure decrease as a result of exaggerated pharmacology induced transient clinical signs and reduced food consumption and body-weight gain. In adult and juvenile dogs, intestine and bone / bone marrow were identified as the main target organs after treatment with selexipag. A delay in the closure of the femoral and/or tibial epiphyseal growth plate was observed in juvenile dogs. A no-observed-adverse-effect level was not established. In juvenile dogs, intussusception due to prostacyclin-related effects on intestinal motility was observed sporadically. Safety margins adapted for IP receptor potency for the active metabolite were 2-fold (based on total exposure) in relation to human therapeutic exposure. The finding did not occur in mouse or rat toxicity studies. Because of the species-specific sensitivity of dogs to develop intussusception, this finding is considered not relevant for adult humans.

Increased bone ossification and related changes in the bone marrow in dog studies are considered to be due to the activation of EP₄ receptors in dogs. As human EP₄ receptors are not activated by selexipag or its active metabolite, this effect is species-specific and, therefore, not relevant to humans.

Selexipag and the active metabolite are not genotoxic on the basis of the overall evidence of conducted genotoxicity studies.

In the 2-year carcinogenicity studies, selexipag caused an increased incidence of thyroid adenomas in mice and Leydig cell adenomas in rats. The mechanisms are rodent-specific. Tortuosity of retinal arterioles was noted after 2 years of treatment only in rats. Mechanistically, the effect is considered to be induced by life-long vasodilation and subsequent changes in ocular haemodynamics. Additional histopathological findings of selexipag were observed only at exposures sufficiently in excess of the maximum human exposure, indicating little relevance to humans.

In a fertility study performed in rats, a prolongation of oestrus cycles resulting in increases in days until copulation was observed at exposures 173-fold above therapeutic exposures (based on total exposures), the no-observed-effect level being 30-fold above therapeutic exposures. Otherwise, fertility parameters were not affected.

Selexipag was not teratogenic in rats and rabbits (exposure margins above therapeutic exposure of 13-fold for selexipag and 43-fold for the active metabolite, based on total exposure). Safety margins for potential IP receptor-related effects on reproduction were 20 for fertility and 5 and 1 (based on free exposure) for embryo-foetal development in rats and rabbits, respectively, when adapted for differences in receptor potency. In the rat pre-/post-natal development study, selexipag induced no effects on maternal and pup reproductive function.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Mannitol (E421),
maize starch,
low substituted hydroxypropyl cellulose,
hydroxypropyl cellulose,
magnesium stearate.

Film coating

Uptravi 200 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide yellow (E172),
carnauba wax.

Uptravi 400 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide red (E172),
carnauba wax.

Uptravi 600 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),

iron oxide red (E172),
iron oxide black (E172),
carnauba wax.

Uptravi 800 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide yellow (E172),
iron oxide black (E172),
carnauba wax.

Uptravi 1,000 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide red (E172),
iron oxide yellow (E172),
carnauba wax.

Uptravi 1,200 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide black (E172),
iron oxide red (E172),
carnauba wax.

Uptravi 1,400 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide yellow (E172),
carnauba wax.

Uptravi 1,600 microgram film-coated tablet

Hypromellose,
propylene glycol,
titanium dioxide (E171),
iron oxide black (E172),
iron oxide red (E172),
iron oxide yellow (E172),
carnauba wax.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

Polyamide / aluminium / HDPE / PE with an embedded desiccant agent / HDPE blister sealed with an aluminium foil.

Uptravi 200 microgram film-coated tablets

Cartons of 10 or 60 film-coated tablets, and 60 or 140 film-coated tablets (titration packs).

Uptravi 400 microgram, 600 microgram, 800 microgram, 1,000 microgram, 1,200 microgram, 1,400 microgram, and 1,600 microgram film-coated tablets

Cartons of 60 film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements for disposal.

7. MARKETING AUTHORISATION HOLDER

Actelion Registration Ltd
Chiswick Tower 13th Floor
389 Chiswick High Road
London W4 4AL
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/15/1083/001
EU/1/15/1083/002
EU/1/15/1083/003
EU/1/15/1083/004
EU/1/15/1083/005
EU/1/15/1083/006
EU/1/15/1083/007
EU/1/15/1083/008
EU/1/15/1083/009
EU/1/15/1083/010
EU/1/15/1083/011

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 12 May 2016

10. DATE OF REVISION OF THE TEXT

June 2017

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.